

Accelerators for Beginners

Rende Steerenberg BE/OP

CERN Accelerator School

Basic Accelerator Science & Technology at CERN

4 – 8 November 2013 – Chavannes de Bogis


- Why Accelerators and Colliders ?
- A very Brief Historic Overview
- The Main Ingredients of an Accelerator

- **Why Accelerators and Colliders ?**
- A very Brief Historic Overview
- The Main Ingredients of an Accelerator

Creating Matter from Energy


$$E = m c^2$$

During the Big Bang Energy was transformed in matter



In our accelerators we provide energy to the particle we accelerate.

In the detectors we observe the matter created



Visible light
 $\lambda = 400 \rightarrow 700 \text{ nm}$




$$\lambda = \frac{hc}{E}$$

X-ray
 $\lambda = 0.01 \rightarrow 10 \text{ nm}$



Increasing the energy will reduce the wavelength

Particle accelerators
 $\lambda < 0.01 \text{ nm}$



Fixed Target



$$E \propto \sqrt{E_{beam}}$$

Much of the energy is lost in the target and only part is used to produce secondary particles

Collider




$$E = E_{beam1} + E_{beam2}$$

All energy will be available for particle production

The Aim

Basics of Accelerator Science & Technology at CERN

Verify the Standard Model



Search for physics beyond the Standard Model

“Standard Model and Beyond” by Paris Sphicas *This afternoon*

- Why Accelerators and Colliders ?
- **A very Brief Historic Overview**
- The Main Ingredients of an Accelerator



Today: ~ **30'000** accelerators operational world-wide*

The **large majority** is used in **industry** and **medicine**

Industrial applications: ~ 20'000*

Medical applications: ~ 10'000*

Les than a **fraction of a percent** is used for **research** and discovery science

Cyclotrons



Synchrotron light sources (e⁻)

Lin. & Circ. accelerators/Colliders

This lecture will concentrate on the CERN type machines of which the majority are **Synchrotrons**

**Source: World Scientific Reviews of Accelerator Science and Technology
A.W. Chao*



- 1932: First accelerator – single passage 160 keV
- Static voltage accelerator
- Limited by the high voltage needed.



Cyclotron


- 1932: 1.2 MeV – 1940: 20 MeV (E.O. Lawrence, M.S. Livingston)
- Constant magnetic field
- Alternating voltage between
- Increasing particle trajectory radius
- Development lead to the synchro-cyclotron to cope with the relativistic effects.

In 1939 Lawrence received the Noble prize for his work.




Betatron

- 1940: Kerst 2.3 MeV and very quickly 300 MeV
- It is actually a transformer with a beam of electrons as secondary winding.
- The magnetic field is used to bend the electrons in a circle, but also to accelerate them.
- A deflecting electrode is used to deflect the particle for extraction.




Linear Accelerator



- Many people involved: Wideroe, Sloan, Lawrence, Alvarez,....
- Main development took place between 1931 and 1946.
- Development was also helped by the progress made on high power high frequency power supplies for radar technology.
- Today still the first stage in many accelerator complexes.
- Limited by energy due to length and single pass.



- 1959: CERN PS and BNL AGS)
- Fixed radius for particle orbit
- Varying magnetic field and radio frequency
- Important focusing of particle beams
- Providing beam for fixed target physics
- Paved the way to colliders




- Why Accelerators and Colliders ?
- A very Brief Historic Overview
- **The Main Ingredients of an Accelerator**

Towards Relativity

Basics of Accelerator Science & Technology at CERN



“Relativity” by Werner Herr This afternoon




In less than 1 hour

“Overview of the CERN Complex” by Paul Collier
 “Sources” by Richard Scrivens
 “LINACS” by Maurizio Vretenar

Wednesday morning


LEIR as an Example



LEIR as an Example

The particle beam:


- arrives through a transfer line
- is injected
- is accelerated over many turns in a “circular” machine
- is extracted
- leaves through a transfer line





Vacuum in a mostly **stainless steel vacuum chamber** is required to **avoid** the particles to **interact** with the **gas molecules**



Especially important for low energy particles and anti-matter particles





In the LHC **vacuum** is also used as **insulator**


“Vacuum Systems” by Vincent Baglin

Thursday afternoon



Injecting & Extracting Particles






“Injection and Extraction” by Wolfgang Bartmann
“Beam Transfer” by Verena Kain
“Kickers and Septa” by Mike Barnes

Wednesday afternoon



Make Particles Circulate



Charged Particles Deviated

Charged Particles are deviated in magnetic fields

Two charged Particles in a homogeneous magnetic field





Lorentz force:

$$F = e v \times B$$

— Particle A
 - - - Particle B

Horizontal motion






Different particles with different initial conditions in a homogeneous magnetic field will cause oscillatory motion in the horizontal plane

The horizontal motion seems to be “stable” ... What about the vertical plane ?

Many particles many initial conditions

Focusing particles, a bit like light



“Transverse Beam Dynamics” by Bernhard Holzer


“Magnets” by Paolo Fessia


“Power Converters” by Jean-Paul Burnet

3 lectures on Tuesday


Thursday morning

Thursday afternoon






Accelerating Beams



~~First attracted
Acceleration
Then again attracted
Deceleration~~

Net result:
No Acceleration

Accelerating Beams




3 lectures on
Tuesday


“Longitudinal Beam Dynamics” by Frank Tecker
“RF Systems” by Erk Jensen

Thursday
morning


The Eyes of Operations



Beam intensity or current measurement



Transverse beam profile/size measurement



Longitudinal beam profile measurements

Measure the LHC luminosity, number of events per surface and time unit.

Any many more beam properties.....

“Beam Instrumentation” by Uli Raich,

Thursday afternoon

Possible Limitations




Machines and elements cannot be built with infinite perfection

Same phase and frequency for driving force and the system can cause resonances



Neighbouring charges with the same polarity experience repelling forces



Moving particles create currents, These currents result in attracting or repelling magnetic fields

Tuesday morning

“Linear Imperfection” by Rogelio
 “Collective effects” Giovanni Rumolo

“Colliders and Beam-Beam” by Tatiana Pieloni

Friday morning



Ever increasing energies and beam intensities, require special techniques

Super conducting magnets, with 8 T or even 11 T instead of 2 T for normal conducting magnets, require cryogenics

High stored beam energies require sophisticated machine protection systems

Thursday morning

“Magnets” by Paolo Fessia


“Cryogenics” by Serge Claudet

“Machine Protection” by Jorg Wenninger

Thursday morning

Friday
afternoon

Everything must be made as simple as possible. But not simpler...



Albert Einstein