


SPACE CHARGE IN CIRCULAR MACHINES

Massimo.Ferrario@LNF.INFN.IT




OUTLINE

- Direct Space Charge Effects
 - The rms emittance concept
 - rms envelope equation
 - Space charge forces
 - Beam (Plasma) emittance oscillations
 - **Intra-beam scattering IBS**
- Image Charge Effects
 - Image self fields
 - Space charge effects in Storage Rings


Space Charge: what does it mean?

The net effect of the **Coulomb** interactions in a multi-particle system can be classified into two regimes:

- 1) **Collisional Regime** ==> dominated by **binary collisions** caused by close particle encounters ==> **Single Particle Effects**



- 2) **Space Charge Regime** ==> dominated by the **self field** produced by the particle distribution, which varies appreciably only over large distances compare to the average separation of the particles ==> **Collective Effects**



A measure for the relative importance of collisional versus collective effects is the:


Debye Length λ_D

Let consider a **non-neutralized** system of **identical charged particles**

We wish to calculate the **effective potential** of a fixed **test charged particle** **surrounded by other particles** that are statistically distributed.



Magnetic focusing




Magnetic focusing

$$\Phi(\vec{r}) = \frac{C}{r}$$

$$C = \frac{e}{4\pi\epsilon_0}$$

$$\Phi_D(\vec{r}) = ?$$

The effective potential of a test charge can be defined as the sum of the potential of the single particle δ and a “perturbation” term Δn .



- $N \Rightarrow$ total number of particles
- $n \Rightarrow$ particle number density (N/V)
- $k_B \Rightarrow$ Boltzmann constant
- $T \Rightarrow$ Temperature
- $k_B T \Rightarrow$ average kinetic energy of the particles

From Poisson Equation:

$$\nabla^2 \Phi_D(\vec{r}) = \frac{e}{\epsilon_0} \delta(\vec{r}) + \frac{e}{\epsilon_0} \Delta n(\vec{r})$$

$$\Delta n = n e^{-e\Phi_D/k_B T} - n \approx -\frac{ne}{k_B T} \Phi_D$$

$$\nabla^2 \Phi_D(\vec{r}) + \lambda_D \Phi_D(\vec{r}) = \frac{e}{\epsilon_0} \delta(\vec{r})$$


$$\lambda_D = \sqrt{\frac{\epsilon_0 k_B T}{e^2 n}}$$

$$\Phi_D(\vec{r}) = \frac{C}{r} e^{-r/\lambda_D}$$

the effective interaction range of the test particle is limited to the **Debye length**

The charges surrounding the test particles have a screening effect

$$\Phi_D(\vec{r}) = \frac{C}{r} e^{-r/\lambda_D} \Rightarrow \begin{cases} \Phi_D(\vec{r}) \approx \Phi(\vec{r}) & \text{for } r \ll \lambda_D \\ \Phi_D(\vec{r}) \ll \Phi(\vec{r}) & \text{for } r \geq \lambda_D \end{cases}$$




$$\Phi_{SC}(\vec{r}) \gg \Phi_D(\vec{r})$$

Smooth functions for the charge and field distributions can be used as long as the Debye length remains small compared to the particle bunch size

The net effect of the **Coulomb** interactions in a multi-particle system can be classified into two regimes:

- 1) **Collisional Regime** ==> dominated by **binary collisions** caused by close particle encounters ==> **Single Particle Effects**


$$\lambda_D = \sqrt{\frac{\epsilon_0 k_B T}{e^2 n}}$$



$$\sigma_{x,y,z} \ll \lambda_D$$

- 2) **Space Charge Regime** ==> dominated by the **self field** produced by the particle distribution, which varies appreciably only over large distances compare to the average separation of the particles ==> **Collective Effects, Single Component Cold Plasma**



$$\sigma_{x,y,z} \gg \lambda_D$$



In a charged particle beam moving at a longitudinal relativistic velocity, assuming that the random transverse motion in the beam is non-relativistic, the Debye length has the following form:


$$\lambda_D = \sqrt{\frac{\epsilon_0 \gamma^2 k_B T}{e^2 n}}$$

$$\langle v_x \rangle = \sqrt{\frac{k_B T}{\gamma m}} \ll c$$




Collisional regime

- 1) **Collisional Regime** ==> dominated by **binary collisions** caused by close particle encounters ==> **Single Particle Effects**




- 1) multiple small-angle scattering events **Intra-Beam Scattering (IBS)**
- 2) large-angle single scattering events **Touschek Effect**

Liouville theorem does not hold anymore under Collisions
=> non Conservative forces involved



The Intrabeam scattering effect

F. Antoniou, Y. Papaphilippou, CERN

- Small angle **multiple Coulomb scattering** effect
 - Redistribution of beam momenta
 - Beam diffusion with impact on the beam quality (Brightness , luminosity, etc)
- **Different approaches** for the probability of scattering
 - Classical Rutherford cross section
 - Quantum approach
 - Relativistic “Golden Rule” for the 2-body scattering process
- **Several theoretical models** and their **approximations** developed over the years
 - Classical models of Piwinski (**P**) and Bjorken-Mtingwa (**BM**)
 - High energy approximations **Bane, CIMP, etc** 
 - Integrals with analytic solutions

- Theoretical models calculate the **IBS growth rates**:

$$\frac{1}{T_i} \propto \frac{N}{\gamma \epsilon_{xn} \epsilon_{yn} \epsilon_{sn}} f(\text{optics}, \gamma, \epsilon_{xn}, \epsilon_{yn}, \epsilon_{sn})$$

- **Complicated integrals** averaged around the rings
 - Depend on **optics** and **beam properties**

- ✓ They have been well benchmarked for hadron machines
- For lepton machines the work is in progress
 - Need to benchmark the IBS effect in the presence of SR and QE
 - Studies and publications from: ATF(2001), CsrTA, SLS, SPEAR3
- Main drawbacks:
 - Gaussian beams assumed
 - Betatron coupling not trivial to be included
 - Impact on damping process (especially in strong IBS regimes)?
- Tracking codes **SIRE** (A. Vivoli) and **CMAD-IBStrack** (M. Pivi, T. Demma)
 - Based on the classical Rutherford cross section

IBS Calculations

Horizontal, vertical and longitudinal **equilibrium states** and **damping times** due to SR damping

The IBS growth rates in one turn (or one time step)

$$\frac{1}{T_i} = \langle f_i \rangle$$

Complicated integrals averaged around the ring.

$$\begin{aligned} \frac{d\varepsilon_x}{dt} &= -\frac{2}{\tau_x} (\varepsilon_x - \varepsilon_{x0}) + \frac{2\varepsilon_x}{T_x(\varepsilon_x, \varepsilon_y, \sigma_p)} \\ \frac{d\varepsilon_y}{dt} &= -\frac{2}{\tau_y} (\varepsilon_y - \varepsilon_{y0}) + \frac{2\varepsilon_y}{T_y(\varepsilon_x, \varepsilon_y, \sigma_p)} \\ \frac{d\sigma_p}{dt} &= -\frac{1}{\tau_p} (\sigma_p - \sigma_{p0}) + \frac{\sigma_p}{T_p(\varepsilon_x, \varepsilon_y, \sigma_p)} \end{aligned}$$

If = 0 **Steady State emittances**

$$\varepsilon'_{x0} = \frac{T_x}{T_x - \tau_x} \varepsilon_{x0}$$

If ≠ 0

- Steady state exists if we are below transition or in the presence of SR
- dt should be much smaller than the IBS growth times
- Good scanning of optics is important in order not to skip large IBS kick points

Observations of IBS emittance growth in the KEK-ATF




FIG. 3. Current dependence of the horizontal emittance: Data for the smallest emittance cases (runs B and D) are shown. The result of a SAD simulation for 0.4% coupling is superimposed.






FIG. 2. Current dependence of the vertical emittance: Data for the smallest emittance cases (runs B and D) are shown. The result of a SAD simulation for 0.4% coupling is superimposed.



Observations of IBS emittance growth in the KEK-ATF




FIG. 4. Current dependence of the momentum spread: Data for the smallest (run D) and the largest emittance case (run E') are shown. (E' means that the data were taken in same condition as E, but on another day.) The results of SAD simulations for 0.4% and 6% coupling are superimposed.






FIG. 5. Current dependence of the bunch length: Data are shown for the runs D', F', and E' (the symbols D', E', and F' indicate that the data were taken for the same condition as D, E, and F, but on another day.) The results of SAD simulations for 0.4%, 6%, and 3% coupling are superimposed.



Beam Thermodynamics

Definition of beam temperature in analogy with kinetic theory of gases :

transverse

$$k_B T_{beam,x} = \gamma m_o \langle v_x^2 \rangle = \gamma m_o \beta^2 c^2 \frac{\epsilon_{rms,x}}{\beta_x}$$

longitudinal

$$k_B T_{beam,z} = \gamma^3 m_o \langle \Delta v_z^2 \rangle = \frac{\beta^2 c^2}{\gamma} m_o \left\langle \left(\frac{\Delta p}{p} \right)^2 \right\rangle$$

$$\Delta v_z = \frac{\beta c}{\gamma^2} \frac{\Delta p}{p}$$

In a **Circular machine** when a particle accelerates above transition energy it becomes slower and behaves like a particle with negative mass:

$$k_B T_{beam,z} = m^* \langle \Delta v_z^2 \rangle = -\frac{\gamma m_o}{\eta} \langle \Delta v_z^2 \rangle$$

$$\eta = \alpha - \frac{1}{\gamma_o^2} = \frac{1}{\gamma_t^2} - \frac{1}{\gamma_o^2}$$

Conservation Law

Let us first consider the ideal machine with a smooth-focusing lattice below transition and negligible dispersion.

The total thermal energy per particle in a smooth linear beam channel is conserved, for a beam with constant energy ($\gamma_0 = \text{const}$)

$$k_B T_x + k_B T_y + k_B T_{\parallel} = \text{const}$$

Coulomb collisions drive the beam toward an isotropic thermal equilibrium, in which case the three temperatures would be the same:

$$k_B T_x = k_B T_y = k_B T_{\parallel} = k_B T_{\text{eq}}$$

We can put the conservation law into the form:

$$\frac{\varepsilon_{rms,x}}{\beta_x} + \frac{\varepsilon_{rms,y}}{\beta_y} + \frac{1}{\gamma_o^2} \left\langle \left(\frac{\Delta p}{p} \right)^2 \right\rangle = \text{const}$$

in a circular machine we must replace $1/\gamma_0$ by:

$$-\eta = \frac{1}{\gamma_0^2} - \alpha = \frac{1}{\gamma_0^2} - \frac{1}{\gamma_t^2}$$

$$\frac{\varepsilon_{rms,x}}{\beta_x} + \frac{\varepsilon_{rms,y}}{\beta_y} - \eta \left\langle \left(\frac{\Delta p}{p} \right)^2 \right\rangle = const$$

This relationship is the invariant for intra-beam scattering derived in 1974 by **Piwinski**. For a circular machine the behavior of the system depends on the sign of η i.e. whether it is **below** transition ($\gamma_0 < \gamma_t$) or **above** ($\gamma_0 > \gamma_t$).

→ **below** transition $\eta < 0$ → **thermal equilibrium can be reached.**

→ **above** transition $\eta > 0$ → **thermal equilibrium is not possible.**

An increase in momentum spread must be balanced by a corresponding increase in the transverse emittances to maintain the “conservation law”

For instance, in the LHC at 7 TeV, although $\gamma = 7461 \gg \gamma_t \approx 53.8$ ($\eta_t \approx 3.4 \times 10^{-4}$), the undesirable growth of the bunch emittances caused by IBS is counterbalanced by the *synchrotron radiation damping* effect.

The growth rate for intra-beam scattering in high-energy circular machines defined as:

$$\frac{1}{\tau_j(\epsilon)} = \frac{1}{\tilde{\epsilon}_j} \frac{d\tilde{\epsilon}_j}{dt}$$

can be written in the relativistic form


$$\frac{1}{\tau_j} = \frac{1}{\tau_0} \langle H_j \rangle = \frac{\pi^2 c r_c^2 m^3 N \ln \Lambda}{\gamma_0 \Gamma} \langle H_j \rangle$$

where N is the total number of particles, Γ the six-dimensional phase-space volume occupied by N , and where the function H_j depends on γ_0 , the emittances $\tilde{\epsilon}_x$, $\tilde{\epsilon}_y$, $\tilde{\epsilon}_z$, and the lattice parameters $\hat{\beta}_x$, D_e , $\hat{\beta}'_x$, D'_e , and $\hat{\beta}_y$. The function H_j is averaged over a lattice period and the subscript j denotes the three orthogonal directions (i.e., $j =$ horizontal (x), vertical (y), and longitudinal (s)).

$$\Gamma_b = (2\pi)^3 \frac{P_0^3}{c^3} \tilde{\epsilon}_x \tilde{\epsilon}_y \tilde{\epsilon}_z = (2\pi)^3 (\beta_0 \gamma_0)^3 m^3 \tilde{\epsilon}_x \tilde{\epsilon}_y \tilde{\epsilon}_z$$


Touschek Effect

In a relativistic storage ring, Coulomb collisions lead to a momentum transfer from the transverse into the longitudinal direction that is amplified by the Lorentz factor γ_0



While the total momentum in the collision is preserved, the two particles emerge from this collision with opposite longitudinal momentum components that are larger by the factor γ_0 than the original transverse momentum component before the collision.

If the longitudinal momentum acquired in such a collision is greater than the momentum acceptance of the rf bucket that keeps the beam longitudinally bunched, the two particles involved in such a collision will be lost.



$$\Delta E_{\max} = -\Delta E_{\min} = 2 \left[\beta_0^3 \gamma_0^3 \frac{\lambda}{2\pi} mc^2 q E_m (\varphi_0 \cos \varphi_0 - \sin \varphi_0) \right]^{1/2}$$

The net result is that the lifetime of the stored beam is reduced.

We do not analyse Touschek scattering in detail, but (as for IBS) simply quote the result. The Touschek lifetime is given by:

$$\frac{1}{\tau} = -\frac{1}{N} \frac{dN}{dt} = \frac{r_e^2 c N}{8\pi\sigma_x\sigma_y\sigma_z} \frac{1}{\gamma^2 \delta_{\max}^3} D\left(\left[\frac{\delta_{\max}\beta_x}{\gamma\sigma_x}\right]^2\right)$$

where N is the number of particles in a bunch, σ_x , σ_y , σ_z are the rms horizontal and vertical beam sizes and bunch length, and δ_{\max} is the energy acceptance of the ring.

Particle loss from the Touschek effect tends to be the dominant limitation on the beam lifetime in low-emittance storage rings, such as those in third-generation synchrotron light sources; and is expected to be the dominant limitation on lifetime in the ILC damping rings.

$$D(\varepsilon) = \sqrt{\varepsilon} \left[-\frac{3}{2} e^{-\varepsilon} + \frac{\varepsilon}{2} \int_{\varepsilon}^{\infty} \frac{\ln u}{u} e^{-u} du + \frac{1}{2} (3\varepsilon - \varepsilon \ln \varepsilon + 2) \int_{\varepsilon}^{\infty} \frac{e^{-u}}{u} du \right]$$

Proceedings of the CAS-CERN Accelerator School: Intensity Limitations in Particle Beams, Geneva, Switzerland, 2–11 November 2015,
edited by W. Herr, CERN Yellow Reports: School Proceedings, Vol. 3/2017, CERN-2017-006-SP (CERN, Geneva, 2017)

Intrabeam Scattering: Anatomy of the Theory


M. Martini

CERN, Geneva, Switzerland

OUTLINE

- Direct Space Charge Effects
 - The rms emittance concept
 - rms envelope equation
 - Space charge forces
 - Beam (Plasma) emittance oscillations
- Image Charge Effects
 - **Image self fields**
 - Space charge effects in Storage Rings

IMAGE SELF FIELDS



Direct self fields





Image self fields




Wake fields

Space Charge

Static Fields: conducting screens


Let us consider a point charge q close to a conducting screen.

The electrostatic field can be derived through the "image method". Since the metallic screen is an equi-potential plane, it can be removed provided that a "virtual" charge is introduced such that the potential is constant at the position of the screen




A constant current in the free space produces circular magnetic field

If $\mu_r \approx 1$, the material, even in the case of a good conductor, does not affect the field lines.



Circular Perfectly Conducting Pipe (Beam at Center)




In the case of cylindrical charge distribution, and $\gamma \rightarrow \infty$, the electric field lines are perpendicular to the direction of motion. The transverse fields intensity can be computed like in the static case, applying the Gauss and Ampere laws.

$$\lambda(r) = \lambda_0 \left(\frac{r}{a} \right)^2; \int_S E_r (2\pi r) \Delta z = \frac{\lambda(r) \Delta z}{\epsilon_0}$$

$$E_r = \frac{\lambda(r)}{2\pi\epsilon_0 r}; B_\theta = \frac{\beta}{c} E_r$$

$$E_r(\mathbf{r}) = \frac{\lambda_0}{2\pi\epsilon_0} \frac{r}{a^2}; B_\theta(\mathbf{r}) = \frac{\lambda_0 \beta}{2\pi\epsilon_0 c} \frac{r}{a^2}$$

$$F_\perp(r) = e(E_r - \beta c B_\theta) = \frac{e}{\gamma^2} E_r$$



there is a cancellation of the electric and magnetic forces



Parallel Plates (beam at center)




In some cases, the beam pipe cross section is such that we can consider only the surfaces closer to the beam, which behave like two parallel plates. In this case, we use the image method to a charge distribution of radius a between two conducting plates $2h$ apart. By applying the superposition principle we get the total image field at a position y inside the beam.

$$E_y^{im}(z, y) = \frac{\lambda(z)}{2\pi \epsilon_0} \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{2nh + y} - \frac{1}{2nh - y} \right]$$

$$E_y^{im}(z, y) = \frac{\lambda(z)}{2\pi \epsilon_0} \sum_{n=1}^{\infty} (-1)^n \frac{-2y}{(2nh)^2 - y^2} \cong \frac{\lambda(z)}{4\pi \epsilon_0 h^2} \frac{\pi^2}{12} y$$

Where we have assumed: $h \gg a > y$.



For d.c. or slowly varying currents, the boundary condition imposed by the conducting plates does not affect the magnetic field. We do not need “image currents” “As a consequence there is no cancellation effect for the fields produced by the “image” charges.

From the divergence equation we derive also the other transverse component, notice the opposite sign:

$$\frac{\partial}{\partial x} E_x^{im} = -\frac{\partial}{\partial y} E_y^{im} \Rightarrow E_x^{im}(z, x) = \frac{-\lambda(z)}{4\pi \epsilon_0 h^2} \frac{\pi^2}{12} x$$

Including also the direct space charge force, we get:


$$\begin{cases} F_x(z, x) = \frac{e\lambda(z)x}{\pi \epsilon_0} \left(\frac{1}{2a^2\gamma^2} - \frac{\pi^2}{48h^2} \right) \\ F_y(z, x) = \frac{e\lambda(z)y}{\pi \epsilon_0} \left(\frac{1}{2a^2\gamma^2} + \frac{\pi^2}{48h^2} \right) \end{cases}$$

Therefore, for $\gamma \gg 1$, and for d.c. or slowly varying currents the cancellation effect applies only for the direct space charge forces. There is no cancellation of the electric and magnetic forces due to the "image" charges.


Time-varying fields

It is necessary to compare the **wall thickness** and the **skin depth** (region of penetration of the e.m. fields) in the conductor.


$$\delta_w \cong \sqrt{\frac{2}{\omega\sigma\mu}}$$



If the **fields penetrate** and pass through the material, we are practically in the **static boundary conditions case**. Conversely, if the **skin depth is very small**, fields do not penetrate, the electric field lines are perpendicular to the wall, as in the static case, while **the magnetic field lines are tangent to the surface**.



Parallel Plates (Beam at Center) a.c. currents




Usually, the frequency beam spectrum is quite rich of harmonics, especially for bunched beams.

It is convenient to decompose the current into a d.c. component, I , for which $\delta_w \gg \Delta_w$, and an a.c. component, \hat{I} , for which $\delta_w \ll \Delta_w$.

While the d.c. component of the magnetic field does not perceive the presence of the material, **its a.c. component is obliged to be tangent at the wall.** For a charge density λ we have $I = \lambda v$.

We can see that this current produces a magnetic field able to cancel the effect of the electrostatic force.



$$\begin{cases} \tilde{E}_y(z, x) = \frac{\tilde{\lambda}(z)y}{\pi \epsilon_0} \frac{\pi^2}{48h^2} \\ \tilde{B}_x(z, x) = \frac{\beta}{c} \tilde{E}_y(z, x) \end{cases}$$

$$\tilde{F}_y(z, x) = e(1 - \beta^2) E_y = \frac{1}{\gamma^2} \frac{e\tilde{\lambda}(z)y}{\pi \epsilon_0} \frac{\pi^2}{48h^2}$$


$$\begin{cases} \tilde{F}_x(z, x) = \frac{e\tilde{\lambda}(z)x}{2\pi \epsilon_0 \gamma^2} \left(\frac{1}{a^2} - \frac{\pi^2}{24h^2} \right) \\ \tilde{F}_y(z, x) = \frac{e\tilde{\lambda}(z)y}{2\pi \epsilon_0 \gamma^2} \left(\frac{1}{a^2} + \frac{\pi^2}{24h^2} \right) \end{cases}$$

There is cancellation of the electric and magnetic forces !!


$$\lambda(z) = \lambda_0 + \tilde{\lambda} \cos(k_z z)$$

D.C.

A.C. ($\delta_w \ll \Delta_w$)



$$F_{\perp}(r) = \frac{e}{\gamma^2} \frac{\lambda(z)}{2\pi \epsilon_0} \frac{r}{a^2}$$




$$F_x(z, x) = \frac{e\lambda_0 x}{\pi \epsilon_0} \left(\frac{1}{2a^2 \gamma^2} - \frac{\pi^2}{48h^2} \right)$$

$$\tilde{F}_x(z, x) = \frac{e\tilde{\lambda}(z)x}{\pi \epsilon_0 \gamma^2} \left(\frac{1}{2a^2} - \frac{\pi^2}{48h^2} \right)$$

$$F_y(z, x) = \frac{e\lambda_0 y}{\pi \epsilon_0} \left(\frac{1}{2a^2 \gamma^2} + \frac{\pi^2}{48h^2} \right)$$

$$\tilde{F}_y(z, x) = \frac{e\tilde{\lambda}(z)y}{\pi \epsilon_0 \gamma^2} \left(\frac{1}{2a^2} + \frac{\pi^2}{48h^2} \right)$$




OUTLINE

- Direct Space Charge Effects
 - The rms emittance concept
 - rms envelope equation
 - Space charge forces
 - Beam (Plasma) emittance oscillations
- Image Charge Effects
 - Image self fields
 - Space charge effects in Storage Rings

Incoherent and Coherent Transverse Effects


When the beam is located at the centre of symmetry of the pipe, the e.m. forces due to space charge and images cannot affect the motion of the centre of mass (**coherent**), but change the trajectory of individual charges in the beam (**incoherent**).



These force may have a complicate dependence on the charge position. A simple analysis is done considering only the linear expansion of the self-fields forces around the equilibrium trajectory.

Self Fields and betatron motion

Consider a perfectly circular accelerator with radius ρ_x . The beam circulates inside the beam pipe. The transverse single particle motion in the linear regime, is derived from the equation of motion. Including the self field forces in the motion equation, we have



$$\frac{d(m\gamma \mathbf{v})}{dt} = \mathbf{F}^{ext}(\vec{r}) + \mathbf{F}^{self}(\vec{r})$$

$$\frac{d\mathbf{v}}{dt} = \frac{\mathbf{F}^{ext}(\vec{r}) + \mathbf{F}^{self}(\vec{r})}{m\gamma}$$

Self Fields and betatron motion

In the analysis of the motion of the particles in presence of the self field, we will adopt a **simplified model where particles execute simple harmonic oscillations around the reference orbit.**

This is the case where the focussing term is constant. Although this condition is never fulfilled in a real accelerator, it provides a reliable model for the description of the beam instabilities

$$x''(s) + K_x x(s) = \frac{1}{\beta^2 E_o} F_x^{self}(x)$$

Q_x , Betatron tune is the n. of betatron oscillations per turn:

$$Q_x = \frac{2\pi\rho_x}{\lambda_\beta} = \frac{2\pi\rho_x\sqrt{K_x}}{2\pi} = \rho_x\sqrt{K_x}$$

$$x''(s) + \left(\frac{Q_x}{\rho_x}\right)^2 x(s) = \frac{1}{\beta^2 E_o} F_x^{self}(x, s)$$

Transverse Incoherent Effects

We take the linear term of the transverse force in the betatron equation:

$$F_x^{s.c.}(x, z) \cong \left(\frac{\partial F_x^{s.c.}}{\partial x} \right)_{x=0} x \qquad x'' + \left(\frac{Q_{x0}}{\rho_x} \right)^2 x = \frac{1}{\beta^2 E_0} \left(\frac{\partial F_x^{s.c.}}{\partial x} \right)_{x=0} x$$

$$x'' + \left(\frac{Q_{x0}}{\rho_x} - \frac{1}{\beta^2 E_0} \left(\frac{\partial F_x^{s.c.}}{\partial x} \right)_{x=0} \right)^2 x = 0$$

$$(Q_{x0} + \Delta Q_x)^2 = Q_{x0}^2 + 2Q_{x0}\Delta Q_x + \cancel{\Delta Q_x^2} \quad \Rightarrow \quad \Delta Q_x = -\frac{\rho_x^2}{2\beta^2 E_0 Q_{x0}} \left(\frac{\partial F_x^{s.c.}}{\partial x} \right)$$

The shift of betatron wave numbers (tune shift) is negative since the space charge forces are defocusing on both planes. Notice that the tune shift is, in general, function of “z”, therefore we have also a tune spread inside the beam. Furthermore, by including higher order terms in the transverse force, we don't have the harmonic oscillator equation any more.

Example: Incoherent betatron tune shift for an **uniform electron beam of radius a , length l_o , inside circular perfectly conducting pipe**

$$\left(\frac{\partial F_x^{s.c.}}{\partial x} \right) = \frac{\partial}{\partial x} \frac{e\lambda_o x}{2\pi\epsilon_o\gamma^2 a^2} = \frac{e\lambda_o}{2\pi\epsilon_o\gamma^2 a^2}$$

$$\Delta Q_x = - \frac{\rho_x^2 N e^2}{4\pi\epsilon_o a^2 \beta^2 \gamma^2 E_o Q_{xo} l_o}$$

$$r_{e,p} = \frac{e^2}{4\pi\epsilon_o m_o c^2} \quad (\text{electrons : } 2.82 \cdot 10^{-15} \text{ m, protons : } 1.53 \cdot 10^{-18} \text{ m})$$

$$\Delta Q_x = - \frac{\rho_x^2 N r_{e,p}}{a^2 \beta^2 \gamma^3 Q_{xo} l_o}$$

For a real bunched beams the space charge forces, and the tune shift depend on the longitudinal and radial position of the charge.

Consequences of the space charge tune shifts


In circular accelerators the values of the betatron tunes should **not be close to rational numbers** in order to avoid the crossing of linear and non-linear resonances where the beam becomes unstable.

The tune spread induced by the space charge force can make hard to satisfy this basic requirement. Typically, in order to avoid major resonances the stability requires


$$|\Delta Q_u| < 0.3$$

Transverse Coherent Effects

If the beam experiences a transverse deflection kick, it starts to perform betatron oscillations as a whole. The beam, source of the space charge fields moves transversely inside the pipe, while individual particles still continue their incoherent motion around the common coherent trajectory.



Circular Perfectly Conducting Pipe (Beam off Center)



$$d = \frac{b^2}{x}$$

The image charge is at a distance “ d ” such that the pipe surface is at constant voltage, and pulls the beam away from the center of the pipe.

The effect is **defocusing**, the horizontal electric image field E and the horizontal force F are:

$$E_{xc}(\mathbf{x}) = \frac{\lambda(z)}{2\pi\epsilon_0} \frac{1}{d-x} \approx \frac{\lambda(z)}{2\pi\epsilon_0} \frac{1}{d} = \frac{\lambda(z)}{2\pi\epsilon_0} \frac{x}{b^2}$$

$$F_{xc}(r) \approx \frac{e\lambda(z)}{2\pi\epsilon_0} \frac{x}{b^2}$$

$$\Delta Q_{xc} = -\frac{\rho_x^2}{2\beta^2 E_0 Q_{x0}} \left(\frac{\partial F_{xc}}{\partial x} \right)$$

$$= -\frac{\rho_x^2}{2\beta^2 E_0 Q_{x0}} \frac{e\lambda(z)}{2\pi\epsilon_0 b^2}$$

$$\Delta Q_{xc} = -\frac{r_e \rho_x^2}{\beta^2 \gamma Q_{x0}} \frac{N}{b^2 l_0}$$

$$r_{e,p} = \frac{e^2}{4\pi\epsilon_0 m_0 c^2}$$

This coherent betatron tune shift, differently from the incoherent one **does not depend on the beam size** but **on the pipe radius** and it is inversely proportional to the beam energy.

Space Charge Effects

CERN-2014-009, pp.331-356

M. Ferrario, M. Migliorati, and L. Palumbo
INFN-LNF and University of Rome “La Sapienza”

