

SYNCHROTRON RADIATION

Lenny Rivkin

Ecole Polytechnique Federale de Lausanne (EPFL)

and Paul Scherrer Institute (PSI), Switzerland

Introduction to Accelerator Physics Course

CERN Accelerator School, Frascati, Italy

2 – 14 November 2008



Useful books and references

A. Hofmann, *The Physics of Synchrotron Radiation*

Cambridge University Press 2004

H. Wiedemann, *Synchrotron Radiation*

Springer-Verlag Berlin Heidelberg 2003

H. Wiedemann, *Particle Accelerator Physics I and II*

Springer Study Edition, 2003

A. W. Chao, M. Tigner, *Handbook of Accelerator Physics and*

Engineering, World Scientific 1999

CERN Accelerator School Proceedings

Synchrotron Radiation and Free Electron Lasers

Grenoble, France, 22 - 27 April 1996

(A. Hofmann's lectures on synchrotron radiation)

CERN Yellow Report 98-04

http://cas.web.cern.ch/cas/CAS_Proceedings-DB.html


Brunnen, Switzerland, 2 – 9 July 2003

CERN Yellow Report 2005-012

<http://cas.web.cern.ch/cas/BRUNNEN/lectures.html>

GENERATION OF SYNCHROTRON RADIATION

Curved orbit of electrons in magnet field




Crab Nebula
6000 light years away



First light observed
1054 AD

GE Synchrotron
New York State



First light observed
1947

Synchrotron radiation: some dates

- 1873 Maxwell's equations
- 1887 Hertz: electromagnetic waves
- 1898 Liénard: retarded potentials
- 1900 Wiechert: retarded potentials
- 1908 Schott: Adams Prize Essay

... waiting for accelerators ...

1940: 2.3 MeV betatron, Kerst, Serber

Maxwell equations (poetry)

*War es ein Gott, der diese Zeichen schrieb
Die mit geheimnisvoll verborg'nem Trieb
Die Kräfte der Natur um mich enthüllen
Und mir das Herz mit stiller Freude füllen.*

Ludwig Boltzman

*Was it a God whose inspiration
Led him to write these fine equations
Nature's fields to me he shows
And so my heart with pleasure glows.*



translated by John P. Blewett

THEORETICAL UNDERSTANDING →

1873 Maxwell's equations

→ made evident that changing charge densities would result in electric fields that would radiate outward

1887 Heinrich Hertz demonstrated such waves:



.... this is of no use whatsoever !

1898 Liénard:

ELECTRIC AND MAGNETIC FIELDS PRODUCED BY A POINT CHARGE MOVING ON AN ARBITRARY PATH

(by means of retarded potentials
...
proposed first by Ludwig Lorenz
in 1867)

L'Éclairage Électrique

REVUE HEBDOMADAIRE D'ÉLECTRICITÉ

DIRECTION SCIENTIFIQUE

A. CORNU, Professeur à l'École Polytechnique, Membre de l'Institut. — A. D'ARSONVAL, Professeur au Collège de France, Membre de l'Institut. — G. LIPPMANN, Professeur à la Sorbonne, Membre de l'Institut. — D. MONNIER, Professeur à l'École centrale des Arts et Manufactures. — H. POINCARÉ, Professeur à la Sorbonne, Membre de l'Institut. — A. POTIER, Professeur à l'École des Mines, Membre de l'Institut. — J. BLONDIN, Professeur agrégé de l'Université.

CHAMP ÉLECTRIQUE ET MAGNÉTIQUE

PRODUIT PAR UNE CHARGE ÉLECTRIQUE CONCENTRÉE EN UN POINT ET ANIMÉE D'UN MOUVEMENT QUELCONQUE

Admettons qu'une masse électrique en mouvement de densité ρ et de vitesse u en chaque point produit le même champ qu'un courant de conduction d'intensité $u\rho$. En conservant les notations d'un précédent article (¹) nous obtiendrons pour déterminer le champ, les équations

$$\frac{1}{4\pi} \left(\frac{d\psi}{dy} - \frac{d^2\psi}{dt^2} \right) = \rho u_x + \frac{df}{dt} \quad (1)$$

$$V^2 \left(\frac{dh}{dy} - \frac{dg}{dt} \right) = - \frac{1}{4\pi} \frac{dx}{dt} \quad (2)$$

avec les analogues déduites par permutation tournante et en outre les suivantes

$$\rho = \left(\frac{df}{dx} + \frac{dg}{dy} + \frac{dh}{dt} \right) \quad (3)$$

$$\frac{dx}{dt} + \frac{dy}{dt} + \frac{dz}{dt} = 0. \quad (4)$$

De ce système d'équations on déduit facilement les relations

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) f = V^2 \frac{dz}{dx} + \frac{d}{dt} (\rho u_x) \quad (5)$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) g = 4\pi V^2 \left[\frac{d}{dt} (\rho u_y) - \frac{d}{dy} (\rho u_z) \right] \quad (6)$$

(¹) La théorie de Lorentz, *L'Éclairage Électrique*, t. XIV, p. 417. a_x, b_y, c_z sont les composantes de la force magnétique et f, g, h , celles du déplacement dans l'éther.

Soient maintenant quatre fonctions ψ, F, G, H définies par les conditions

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) \psi = - 4\pi V^2 \rho. \quad (7)$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) F = - 4\pi V^2 \rho u_x \quad (8)$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) G = - 4\pi \rho u_y \quad (9)$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) H = - 4\pi V^2 \rho u_z \quad (10)$$

On satisfara aux conditions (5) et (6) en prenant

$$4\pi f = - \frac{d\psi}{dx} - \frac{1}{V^2} \frac{dF}{dt} \quad (11)$$

$$x = \frac{dH}{dy} - \frac{dG}{dt}. \quad (12)$$

Quant aux équations (1) à (4), pour qu'elles soient satisfaites, il faudra que, en plus de (7) et (8), on ait la condition

$$\frac{d\psi}{dt} + \frac{dF}{dx} + \frac{dG}{dy} + \frac{dH}{dt} = 0. \quad (13)$$

Occupons-nous d'abord de l'équation (7). On sait que la solution la plus générale est la suivante :

$$\psi = \int \frac{\rho [x', y', z', t - \frac{r}{V}]}{r} du' \quad (14)$$

Fig. 1. First page of Liénard's 1898 paper.

1912 Schott:

COMPLETE THEORY OF
SYNCHROTRON RADIATION
IN ALL THE GORY DETAILS
(327 pages long)

... to be forgotten for 30 years
(on the usefulness of prizes)

ELECTROMAGNETIC RADIATION

AND THE MECHANICAL REACTIONS
ARISING FROM IT

BEING AN ADAMS PRIZE ESSAY IN THE
UNIVERSITY OF CAMBRIDGE

by

G. A. SCHOTT, B.A., D.Sc.

Professor of Applied Mathematics in the University College of Wales, Aberystwyth
Formerly Scholar of Trinity College, Cambridge

Cambridge :
at the University Press

1912

Donald Kerst: first betatron (1940)



*"Ausserordentlich hochgeschwindigkeitelektronen
entwickelndenschwerarbeitsbeigollitron"*

Synchrotron radiation: some dates


- 1946 Blewett observes **energy loss**
due to synchrotron radiation
100 MeV betatron
- 1947 First **visual** observation of SR
70 MeV synchrotron, GE Lab
- 1949 **Schwinger PhysRev paper**
- ...
- 1976 **Madey: first demonstration of**
Free Electron laser

Crab Nebula
6000 light years away



First light observed
1054 AD


GE Synchrotron
New York State



First light observed
1947


Why do they radiate?

Charge at rest: Coulomb field, no radiation




Uniformly moving charge
does not radiate


But! Cerenkov!



Accelerated charge



Bremsstrahlung or breaking radiation



Liénard-Wiechert potentials

$$\varphi(t) = \frac{1}{4\pi\epsilon_0} \frac{q}{[\mathbf{r}(1 - \mathbf{n} \cdot \vec{\beta})]_{ret}}$$

$$\vec{\mathbf{A}}(t) = \frac{q}{4\pi\epsilon_0 c^2} \left[\frac{\vec{\mathbf{v}}}{[\mathbf{r}(1 - \mathbf{n} \cdot \vec{\beta})]_{ret}} \right]$$

and the electromagnetic fields:

$$\nabla \cdot \vec{\mathbf{A}} + \frac{1}{c^2} \frac{\partial \varphi}{\partial t} = 0 \quad (\text{Lorentz gauge})$$

$$\vec{\mathbf{B}} = \nabla \times \vec{\mathbf{A}}$$

$$\vec{\mathbf{E}} = -\nabla \varphi - \frac{\partial \vec{\mathbf{A}}}{\partial t}$$


Fields of a moving charge

$$\vec{E}(t) = \frac{q}{4\pi\epsilon_0} \left[\frac{\vec{n} - \vec{\beta}}{(1 - \vec{n} \cdot \vec{\beta})^3 \gamma^2} \cdot \frac{1}{\mathbf{r}^2} \right]_{ret} +$$


$$\frac{q}{4\pi\epsilon_0 c} \left[\frac{\vec{n} \times [(\vec{n} - \vec{\beta}) \times \vec{\dot{\beta}}]}{(1 - \vec{n} \cdot \vec{\beta})^3 \gamma^2} \cdot \frac{1}{\mathbf{r}} \right]_{ret}$$

$$\vec{B}(t) = \frac{1}{c} [\vec{n} \times \vec{E}]$$


Transverse acceleration



Radiation field quickly separates itself from the Coulomb field




Longitudinal acceleration




Radiation field cannot separate itself from the Coulomb field

Moving Source of Waves



Time compression

Electron with velocity β emits a wave with period T_{emit} while the observer sees a different period T_{obs} because the electron was moving towards the observer



$$T_{\text{obs}} = (1 - \mathbf{n} \cdot \boldsymbol{\beta}) T_{\text{emit}}$$

The wavelength is shortened by the same factor

$$\lambda_{\text{obs}} = (1 - \beta \cos \theta) \lambda_{\text{emit}}$$


in ultra-relativistic case, looking along a tangent to the trajectory

$$\lambda_{\text{obs}} = \frac{1}{2\gamma^2} \lambda_{\text{emit}}$$


since

$$1 - \beta = \frac{1 - \beta^2}{1 + \beta} \approx \frac{1}{2\gamma^2}$$

Radiation is emitted into a narrow cone



$$\theta = \frac{1}{\gamma} \cdot \theta_e$$






V << C

V ≈ C

Sound waves (non-relativistic)

Angular collimation



$$\theta = \frac{v_{s\perp}}{v_{s\parallel} + v} = \frac{v_{s\perp}}{v_{s\parallel}} \cdot \frac{1}{1 + \frac{v}{v_s}} \approx \theta_e \cdot \frac{1}{1 + \frac{v}{v_s}}$$

Doppler effect (moving source of sound)

$$\lambda_{heard} = \lambda_{emitted} \left(1 - \frac{v}{v_s} \right)$$

Synchrotron radiation power

Power emitted is proportional to:

$$P \propto E^2 B^2$$

$$P_\gamma = \frac{c C_\gamma}{2\pi} \cdot \frac{E^4}{\rho^2}$$

$$C_\gamma = \frac{4\pi}{3} \frac{r_e}{(m_e c^2)^3} = 8.858 \cdot 10^{-5} \left[\frac{\text{m}}{\text{GeV}^3} \right]$$

The power is all too real!




fig. 12. Damaged X-ray ring front end gate valve. The power incident on the valve was approximately 1 kW for a duration estimated to 2–10 min and drilled a hole through the valve plate.

Synchrotron radiation power

Power emitted is proportional to:

$$P \propto E^2 B^2$$

$$P_\gamma = \frac{c C_\gamma}{2\pi} \cdot \frac{E^4}{\rho^2}$$

$$P_\gamma = \frac{2}{3} \alpha \hbar c^2 \cdot \frac{\gamma^4}{\rho^2}$$

$$C_\gamma = \frac{4\pi}{3} \frac{r_e}{(m_e c^2)^3} = 8.858 \cdot 10^{-5} \left[\frac{\text{m}}{\text{GeV}^3} \right]$$

$$\alpha = \frac{1}{137}$$

Energy loss per turn:


$$\hbar c = 197 \text{ Mev} \cdot \text{fm}$$

$$U_0 = C_\gamma \cdot \frac{E^4}{\rho}$$

$$U_0 = \frac{4\pi}{3} \alpha \hbar c \frac{\gamma^4}{\rho}$$

Typical frequency of synchrotron light

Due to extreme collimation of light observer sees only a small portion of electron trajectory (**a few mm**)



Pulse length:
difference in times it takes an electron and a photon to cover this distance


$$\Delta t \sim \frac{l}{\beta c} - \frac{l}{c} = \frac{l}{\beta c}(1 - \beta)$$

$$\omega \sim \frac{1}{\Delta t} \sim \gamma^3 \omega_0$$

$$\Delta t \sim \frac{2\rho}{\gamma c} \cdot \frac{1}{2\gamma^2}$$

Spectrum of synchrotron radiation

- Synchrotron light comes in a series of flashes every T_0 (revolution period)



- the spectrum consists of harmonics of

$$\omega_0 = \frac{1}{T_0}$$

- flashes are extremely short: harmonics reach up to very high frequencies

$$\omega_{typ} \equiv \gamma^3 \omega_0$$

- At high frequencies the individual harmonics overlap

$$\begin{aligned}\omega_0 &\sim 1 \text{ MHz} \\ \gamma &\sim 4000 \\ \omega_{typ} &\sim 10^{16} \text{ Hz !}\end{aligned}$$


continuous spectrum !

$$\frac{dP}{d\omega} = \frac{P_{\text{tot}}}{\omega_c} S\left(\frac{\omega}{\omega_c}\right)$$

$$S(x) = \frac{9\sqrt{3}}{8\pi} x \int_x^\infty K_{5/3}(x') dx' \quad \int_0^\infty S(x') dx' = 1$$

$$P_{\text{tot}} = \frac{2}{3} \hbar c^2 \alpha \frac{\gamma^4}{\rho^2}$$

$$\omega_c = \frac{3c\gamma^3}{2\rho}$$



A useful approximation

Spectral flux from a dipole magnet with field B


$$\text{Flux} \left[\frac{\text{photons}}{\text{s} \cdot \text{mrad} \cdot 0.1\% \text{BW}} \right] = 2.46 \cdot 10^{13} E[\text{GeV}] I[\text{A}] G_1(x)$$

Approximation: $G_1 \approx A x^{1/3} g(x)$

$$g(x) = \left[\left(1 - \left(\frac{x}{x_L} \right)^N \right)^{\frac{1}{S}} \right]$$


$$A = 2.11, N = 0.848$$

$$x_L = 28.17, S = 0.0513$$



Werner Joho, PSI

Synchrotron radiation flux for different electron energies



Angular divergence of radiation

The rms opening angle R'

- at the critical frequency:

$$\omega = \omega_c \quad R' \approx \frac{0.54}{\gamma}$$

- well below

$$\omega \ll \omega_c \quad R' \approx \frac{1}{\gamma} \left(\frac{\omega_c}{\omega} \right)^{1/3} \approx 0.4 \left(\frac{\lambda}{\rho} \right)^{1/3}$$


independent of γ !

- well above


$$\omega \gg \omega_c \quad R' \approx \frac{0.6}{\gamma} \left(\frac{\omega_c}{\omega} \right)^{1/2}$$

Angular divergence of radiation

- at the critical frequency




- well below



$$\omega = 0.2 \omega_c$$

- well above


$$\omega = 2 \omega_c$$





$$\gamma\theta$$

Polarisation

Synchrotron radiation observed in the plane of the particle orbit is horizontally polarized, i.e. the electric field vector is horizontal




Observed out of the horizontal plane, the radiation is elliptical




Seeing the electron beam (SLS)

X rays



$$\sigma_x \sim 55 \mu\text{m}$$

visible light, vertically polarised



END