


Introduction to Transverse Beam Optics II


Bernhard Holzer, DESY-HERA



Reminder: the ideal world

$$\begin{pmatrix} x \\ x' \end{pmatrix}_s = M \cdot \begin{pmatrix} x \\ x' \end{pmatrix}_0$$

$$M_{foc} = \begin{pmatrix} \cos(\sqrt{|K|}s) & \frac{1}{\sqrt{|K|}} \sin(\sqrt{|K|}s) \\ -\sqrt{|K|} \sin(\sqrt{|K|}s) & \cos(\sqrt{|K|}s) \end{pmatrix}_0$$



The Beta Function

Beam parameters of a typical high energy ring:
I_p = 100 mA
 particles per bunch: *N* ≈ 10¹¹




Example: HERA Bunch pattern

... question: do we really have to calculate some 10¹¹ single particle trajectories ?


$$x(s) = \sqrt{\epsilon} \sqrt{\beta(s)} \cdot \cos(\psi(s) + \phi)$$

$$\psi(s) = \int_0^s \frac{ds}{\beta(s)}$$




Beam Emittance and Phase Space Ellipse

$$\varepsilon = \gamma(s) * x^2(s) + 2\alpha(s)x(s)x'(s) + \beta(s)x'(s)^2$$



Usually we get in a quadrupole $\alpha(s) = 0$

*Inside foc quadrupoles β reaches maximum
→ largest aperture needed*



... the not so ideal world

1.) Emittance ... so sorry $\mathcal{E} \neq \text{const.}$

According to Hamiltonian mechanics:
phase space diagram relates the variables q and p

$q = \text{position} = x$
 $p = \text{momentum} = mc\gamma\beta$

Liouville's Theorem: $\int p dq = \text{const}$

for convenience (i.e. *because we are lazy bones*) we use in accelerator theory:

$$x' = \frac{dx}{ds} = \frac{dx}{dt} \frac{dt}{ds} = \frac{\beta_x}{\beta} \quad \text{where } \beta = v/c$$

$$\int p dq = \text{const} = mc \int \gamma \beta_x dx = mc \gamma \beta \int x' dx$$

$$\Rightarrow \mathcal{E} = \int x' dx \propto \frac{1}{\beta\gamma}$$

*the beam emittance shrinks
during acceleration $\mathcal{E} \sim 1/\gamma$*

2.) Dispersion

Momentum error: $\frac{\Delta p}{p} \neq 0$

Question: do you remember yesterday on page 11 ... sure you do:

Force acting on the particles

$$F = m \frac{d^2}{dt^2} (x + \rho) - \frac{mv^2}{x + \rho} = eB_z v$$

$$x'' - \frac{1}{\rho} \left(1 - \frac{x}{\rho}\right) = \frac{eB_0}{mv} + \frac{exg}{mv} \rightarrow \frac{1}{mv} = \frac{1}{p_0 + \Delta p} \approx \frac{1}{p_0} \left(1 - \frac{\Delta p}{p_0}\right)$$

neglecting higher order terms ...

$$x'' + x \left(\frac{1}{\rho^2} - k \right) = \frac{\Delta p}{p} \cdot \frac{1}{\rho}$$

Momentum spread of the beam adds a term on the r.h.s. of the equation of motion.
→ **inhomogeneous differential equation.**

$$x'' + x\left(\frac{1}{\rho^2} - k\right) = \frac{\Delta p}{p} \cdot \frac{1}{\rho}$$

general solution:

$$x(s) = x_h(s) + x_i(s)$$

$$\begin{cases} x_h''(s) + K(s) \cdot x_h(s) = 0 \\ x_i''(s) + K(s) \cdot x_i(s) = \frac{1}{\rho} \cdot \frac{\Delta p}{p} \end{cases}$$

Normalise with respect to $\Delta p/p$:


$$D(s) = \frac{x_i(s)}{\frac{\Delta p}{p}}$$

Dispersion function $D(s)$

- * *is that special orbit, an ideal particle would have for $\Delta p/p = 1$*
- * *the orbit of any particle is the sum of the well known x_β and the dispersion*
- * *as $D(s)$ is just another orbit it will be subject to the focusing properties of the lattice*

Dispersion

Example: uniform dipole field



Matrix formalism:

$$x(s) = x_\beta(s) + D(s) \cdot \frac{\Delta p}{p}$$

$$x(s) = C(s) \cdot x_0 + S(s) \cdot x'_0 + D(s) \cdot \frac{\Delta p}{p}$$

$$\begin{pmatrix} x \\ x' \end{pmatrix}_s = \begin{pmatrix} C & S \\ C' & S' \end{pmatrix} \begin{pmatrix} x \\ x' \end{pmatrix}_0 + \frac{\Delta p}{p} \begin{pmatrix} D \\ D' \end{pmatrix}$$

$$\begin{pmatrix} x \\ x' \\ \Delta p/p \end{pmatrix}_s = \begin{pmatrix} C & S & D \\ C' & S' & D' \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ x' \\ \Delta p/p \end{pmatrix}_0$$

Example HERA

$$x_\beta = 1 \dots 2 \text{ mm}$$

$$D(s) \approx 1 \dots 2 \text{ m}$$


$$\Delta p/p \approx 1 \cdot 10^{-3}$$

Amplitude of Orbit oscillation

contribution due to Dispersion \approx beam size

Calculate D, D'

$$D(s) = S(s) \int_{s_0}^{s_1} \frac{1}{\rho} C(\tilde{s}) d\tilde{s} - C(s) \int_{s_0}^{s_1} \frac{1}{\rho} S(\tilde{s}) d\tilde{s}$$



Example: Drift

$$M_{Drift} = \begin{pmatrix} 1 & l \\ 0 & 1 \end{pmatrix}$$

$$M_{Drift} = \begin{pmatrix} 1 & l & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$D(s) = S(s) \underbrace{\int_{s_0}^{s_1} \frac{1}{\rho} C(\tilde{s}) d\tilde{s}}_{=0} - C(s) \underbrace{\int_{s_0}^{s_1} \frac{1}{\rho} S(\tilde{s}) d\tilde{s}}_{=0}$$

Example: Dipole

$$M_{Dipole} = \begin{pmatrix} \cos \frac{l}{\rho} & \rho \sin \frac{l}{\rho} \\ -\frac{1}{\rho} \sin \frac{l}{\rho} & \cos \frac{l}{\rho} \end{pmatrix}$$

→

$$D(s) = \rho \cdot \left(1 - \cos \frac{l}{\rho}\right)$$

$$D'(s) = \sin \frac{l}{\rho}$$

3.) Momentum Compaction Factor:


The dispersion function relates the momentum error of a particle to the horizontal orbit coordinate.

inhomogeneous differential equation

$$x'' + K(s) * x = \frac{1}{\rho} \frac{\Delta p}{p}$$

general solution

$$x(s) = x_{\beta}(s) + D(s) \frac{\Delta p}{p}$$




But it does much more:

it changes the length of the off - energy - orbit !!

*particle with a displacement x to the design orbit
 → path length dl ...*

$$\frac{dl}{ds} = \frac{\rho + x}{\rho}$$

$$\rightarrow dl = \left(1 + \frac{x}{\rho(s)} \right) ds$$



circumference of an off-energy closed orbit

$$l_{\Delta E} = \oint dl = \oint \left(1 + \frac{x_{\Delta E}}{\rho(s)} \right) ds$$

remember:

$$x_{\Delta E}(s) = D(s) \frac{\Delta p}{p}$$

$$\delta l_{\Delta E} = \frac{\Delta p}{p} \oint \left(\frac{D(s)}{\rho(s)} \right) ds$$

** The lengthening of the orbit for off-momentum particles is given by the dispersion function and the bending radius.*

Definition:
$$\frac{\delta l_\varepsilon}{L} = \alpha_{cp} \frac{\Delta p}{p}$$

$$\rightarrow \alpha_{cp} = \frac{1}{L} \oint \left(\frac{D(s)}{\rho(s)} \right) ds$$

For first estimates assume:
$$\frac{1}{\rho} = \text{const}$$

$$\int_{\text{dipoles}} D(s) ds = l_{\text{dipoles}} \cdot \langle D \rangle_{\text{dipole}}$$

$$\alpha_{cp} = \frac{1}{L} l_{\text{dipoles}} \langle D \rangle \frac{1}{\rho} = \frac{1}{L} 2\pi\rho \langle D \rangle \frac{1}{\rho} \rightarrow \alpha_{cp} \approx \frac{2\pi}{L} \langle D \rangle \approx \frac{\langle D \rangle}{R}$$


Assume: $v \approx c$

$$\rightarrow \frac{\delta T}{T} = \frac{\delta l_\varepsilon}{L} = \alpha_{cp} \frac{\Delta p}{p}$$

α_{cp} combines via the dispersion function the momentum spread with the longitudinal motion of the particle.

Tune and Quadrupoles

Question: *what will happen, if you do not make too many mistakes and your **particle performs one complete turn** ?*



Transfer Matrix from point „0“ in the lattice to point „s“:

$$M = \begin{pmatrix} \sqrt{\frac{\beta_s}{\beta_0}} (\cos \psi_s + \alpha_0 \sin \psi_s) & \sqrt{\beta_s \beta_0} \sin \psi_s \\ \frac{(\alpha_0 - \alpha_s) \cos \psi_s - (1 + \alpha_0 \alpha_s) \sin \psi_s}{\sqrt{\beta_s \beta_0}} & \sqrt{\frac{\beta_0}{\beta_s}} (\cos \psi_s - \alpha_s \sin \psi_s) \end{pmatrix}$$

Matrix for one complete turn

the Twiss parameters are periodic in L:

$$\beta(s + L) = \beta(s)$$


$$\alpha(s + L) = \alpha(s)$$

$$\gamma(s + L) = \gamma(s)$$

$$M_{\text{turn}} = \begin{pmatrix} C & S \\ C' & S' \end{pmatrix} = \begin{pmatrix} \cos \psi_{\text{turn}} + \alpha \sin \psi_{\text{turn}} & \beta \sin \psi_{\text{turn}} \\ -\gamma \sin \psi_{\text{turn}} & \cos \psi_{\text{turn}} - \alpha \sin \psi_{\text{turn}} \end{pmatrix}$$

Definition: phase advance of the particle oscillation per revolution in units of 2π is called tune

$$Q = \frac{\Delta \psi_{\text{turn}}}{2\pi} = \frac{\mu}{2\pi}$$



Quadrupole Error in the Lattice

optic *perturbation* described by *thin lens quadrupole*

$$M_{dist} = M_{\Delta k} \cdot M_0 = \underbrace{\begin{pmatrix} 1 & 0 \\ -\Delta k ds & 1 \end{pmatrix}}_{\text{quad error}} \cdot \underbrace{\begin{pmatrix} \cos \psi_{turn} + \alpha \sin \psi_{turn} & \beta_0 \sin \psi_{turn} \\ -\gamma \sin \psi_{turn} & \cos \psi_{turn} - \alpha \sin \psi_{turn} \end{pmatrix}}_{\text{ideal storage ring}}$$

$$M_{dist} = \begin{pmatrix} \cos \psi + \alpha \sin \psi & \beta \sin \psi \\ \Delta k ds \cdot (\cos \psi + \alpha \sin \psi) - \gamma \sin \psi & -\Delta k ds \cdot \beta \sin \psi + \cos \psi - \alpha \sin \psi \end{pmatrix}$$

rule for getting the tune

$$\text{Trace}(M) = 2 \cos \psi = 2 \cos \psi_0 + \Delta k ds \beta \sin \psi_0$$

$$\psi = \psi_0 + \Delta \psi \quad \text{Quadrupole error} \rightarrow \text{Tune Shift}$$

$$\cos(\psi_0 + \Delta\psi) = \cos\psi_0 + \frac{\Delta k ds \beta \sin\psi_0}{2}$$

remember the old fashioned trigonometric stuff and assume that the error is small !!!

$$\underbrace{\cos\psi_0 \cdot \cos\Delta\psi}_{\approx 1} - \underbrace{\sin\psi_0 \cdot \sin\Delta\psi}_{\approx \Delta\psi} = \cos\psi_0 + \frac{\Delta k ds \beta \sin\psi_0}{2}$$

$$\Delta\psi = \frac{\Delta k ds \beta}{2}$$

and referring to Q instead of ψ : $\psi = 2\pi Q$


$$\Delta Q = \int_{s_0}^{s_0+L} \frac{\Delta k(s) \beta(s) ds}{4\pi}$$

a quadrupole error leads to a shift of the tune:


$$\Delta Q = \int_{s_0}^{s_0+l} \frac{\Delta k \beta(s)}{4\pi} ds \approx \frac{\Delta k l_{quad} \bar{\beta}}{4\pi}$$

- ! the tune shift is **proportional to the β -function** at the quadrupole
- !! field quality, power supply tolerances etc are much tighter at places where β is large
- !!! mini beta quads: $\beta \approx 1900$
arc quads: $\beta \approx 80$
- !!!! β is a measure for the sensitivity of the beam

Example: measurement of β in a storage ring:



tune spectrum ...




tune shift as a function of a gradient change

Chromaticity: ξ

Influence of external fields on the beam: *prop. to magn. field & prop. zu $1/p$*

dipole magnet

$$\alpha = \frac{\int B^* dl}{p/e}$$



$$x_D = D(s) * \frac{\Delta p}{p}$$

focusing lens

$$k = \frac{g}{p/e}$$




Figure 29: FODO cell

particle having ...
to high energy
to low energy
ideal energy

Chromaticity: ξ

$$k = \frac{g}{\frac{p}{e}}$$

$$p = p_0 + \Delta p$$

in case of a momentum spread:

$$k = \frac{e \cdot g}{p_0 + \Delta p} \approx \frac{e}{p_0} \left(1 - \frac{\Delta p}{p_0}\right) \cdot g = k_0 - \Delta k$$

$$\Delta k = \frac{\Delta p}{p_0} \cdot k_0$$

... which acts like a quadrupole error in the machine and leads to a tune spread:

$$dQ = \frac{\Delta p}{p_0} \cdot \frac{1}{4\pi} k_0 \cdot \beta(s) ds$$

definition of chromaticity:

$$\Delta Q = \xi \cdot \frac{\Delta p}{p_0}$$

Problem: chromaticity is generated by the lattice itself !!

ξ is a **number** indicating the **size of the tune spot** in the working diagram,

ξ is always created if the beam is focussed

→ it is determined by the focusing strength **k** of all quadrupoles

$$\xi := \frac{1}{4\pi} * \oint k(s)\beta(s)ds$$

k = quadrupole strength

β = **betafunction** indicates the beam size ... and even more the **sensitivity of the beam to external fields**

Example: HERA

HERA-p: $\xi = -70 \dots -80$
 $\Delta p/p = 0.5 * 10^{-3}$
 $Q = 0.257 \dots 0.337$

→ **Some particles get very close to resonances and are lost**

Correction of ξ :

1.) sort the particles according to their momentum

$$x_D(s) = D(s) \frac{\Delta p}{p}$$

2.) apply a magnetic field that rises quadratically with x (sextupole field)

$$B_x = \tilde{g}xz$$



$$B_z = \frac{1}{2} \tilde{g}(x^2 - z^2)$$



$$\frac{\partial B_x}{\partial z} = \frac{\partial B_z}{\partial x} = \tilde{g}$$

linear rising
„gradient“:

Sextupole Magnets:



normalised quadrupole strength:

$$k_{sext} = \frac{\tilde{g}x}{p/e} = m_{sext} \cdot x$$


$$k_{sext} = m_{sext} \cdot D \frac{\Delta p}{p}$$

corrected chromaticity:

$$\xi = \frac{-1}{4\pi} \oint \{k(s) - mD(s)\} \beta(s) ds$$

Chromaticity in the FoDo Lattice

$$\xi = -\frac{1}{4\pi} \int \beta(s) * k(s) ds$$



β -Function in a FoDo

$$\hat{\beta} = \frac{(1 + \sin \frac{\mu}{2})L}{\sin \mu}$$

$$\check{\beta} = \frac{(1 - \sin \frac{\mu}{2})L}{\sin \mu}$$

$$\xi \approx -\frac{1}{4\pi} N * \frac{\hat{\beta} - \check{\beta}}{f_Q}$$

$$\xi = -\frac{1}{4\pi} N * \frac{1}{f_Q} * \left\{ \frac{L(1 + \sin \frac{\mu}{2}) - L(1 - \sin \frac{\mu}{2})}{\sin \mu} \right\}$$

using some *TLC transformations* ... ξ can be expressed in a very simple form:

$$\xi = -\frac{1}{4\pi} N \frac{1}{f_Q} \frac{2L \sin \frac{\mu}{2}}{\sin \mu}$$

$$\xi = -\frac{1}{4\pi} N \frac{1}{f_Q} \frac{L \sin \frac{\mu}{2}}{\sin \frac{\mu}{2} \cos \frac{\mu}{2}}$$

remember ...

$$\sin x = 2 \sin \frac{x}{2} \cos \frac{x}{2}$$

$$\xi_{Cell} = -\frac{1}{4\pi f_Q} * \frac{L \tan \frac{\mu}{2}}{\sin \frac{\mu}{2}}$$

putting ...

$$\sin \frac{\mu}{2} = \frac{L}{4f_Q}$$


$$\xi_{Cell} = -\frac{1}{\pi} * \tan \frac{\mu}{2}$$

contribution of one FoDo Cell to the chromaticity of the ring:

Chromaticity

$$\xi = \frac{-1}{4\pi} \oint k(s) \beta(s) ds$$

question: main contribution to ξ in a lattice ... ?



Resume':

beam emittance

$$\varepsilon \propto \frac{1}{\beta\gamma}$$

dispersion orbit

$$x(s) = x_\beta(s) + D(s) \frac{\Delta p}{p}$$

momentum compaction

$$\frac{\delta l_\varepsilon}{L} = \alpha_{cp} \frac{\Delta p}{p}$$

$$\alpha_{cp} \approx \frac{2\pi}{L} \langle D \rangle \approx \frac{\langle D \rangle}{R}$$

quadrupole error

$$\Delta Q = \int_{s_0}^{s_0+l} \frac{\Delta k(s) \beta(s) ds}{4\pi}$$

chromaticity

$$\xi := \frac{1}{4\pi} * \oint k(s) \beta(s) ds$$

Question:

... after all the very exciting question:

why do our particles not obey gravity and just fall down in the storage ring ????

Answer will be discussed in the evening, having a good glass of red wine at the bar.